

(June 15, 2017)

# **Mirror-Visual Feedback Recent Reviews:**

## **A list of placebo-controlled clinical trials and case reports**

**(A lay summary of our work from the BBC World Review for  
non-specialists, patients, and physicians is included at the end.)**



Controlled Clinical Trials using Mirror Visual Feedback (MVF) have now established the efficacy of the procedure for many neurological conditions. There are over 100 published studies.

## **Evidence for the Utility of MVF in Neurology**

Recent work is highlighted in red

### **Mirror Visual Feedback (MVF) for Stroke**

1. Altschuler EL, Wisdom SB, Stone L, Foster C, Galasko D, Llewellyn DME, Ramachandran VS (1999) et al. Rehabilitation of hemiparesis after stroke with a mirror. *Lancet* 1999;353:2035-6
2. The Centers for Disease Control and Prevention. (2015). Stroke. Retrieved from <http://www.cdc.gov/stroke/index.htm>
3. Sathian K, Greenspan AI, Wolf SL. Doing it with mirrors: a case study of a novel approach to neurorehabilitation. *Neurorehabil Neural Repair* 2000;14:73-6.
4. Stevens JA, Stoykov ME. Using motor imagery in the rehabilitation of hemiparesis. *Arch Phys Med Rehabil* 2003;84:1090-2.
5. Stevens JA, Stoykov ME. Simulation of bilateral movement training through mirror reflection: a case report demonstrating an occupational therapy technique for hemiparesis. *Top Stroke Rehabil* 2004;11:59-66
6. Sütbeyaz S, Yavuzer G, Sezer N, Koseoglu BF . Mirror therapy enhances lower-extremity motor recovery and motor functioning after stroke: a randomized controlled trial. *Arch Phys Med Rehabil* 2007;88:555-9.
7. Altschuler EL, Hu J. Mirror therapy in a patient with a fractured wrist and no active wrist extension. *Scand J Plast Reconstr Surg Hand Surg* 2008;42:110-1.
8. Ramachandran VS, Altschuler EL. The use of visual feedback, in particular mirror visual feedback, in restoring brain function. *Brain*. 2009 Jul;132(Pt 7):1693-710.

9. Lundquist CB, Nielsen JF. Left/right judgement does not influence the effect of mirror therapy after stroke. *Disabil Rehabil.* 2014;36(17):1452-6.
10. Nilsen DM, DiRusso T. Using mirror therapy in the home environment: a case report. *Am J Occup Ther.* 2014 May-Jun;68(3):e84-9.
11. Ramachandran VS, Altschuler EL, Hillyer S. Mirror agnosia. *Proc Biol Sci.* 1997 May 22;264(1382):645-7.
12. Ramachandran VS1, Altschuler EL, Stone L, Al-Aboudi M, Schwartz E, Siva N. Can mirrors alleviate visual hemineglect? *Med Hypotheses.* 1999 Apr;52(4):303-5.
13. Schuster-Amft C, Henneke A, Hartog-Keisker B, Holper L, Siekierka E, Chevrier E, Pyk P, Kollias S, Kiper D, Eng K. Intensive virtual reality-based training for upper limb motor function in chronic stroke: a feasibility study using a single case experimental design and fMRI. *Disabil Rehabil Assist Technol.* 2015;10(5):385-92.
14. Shiri S, Feintuch U, Lorber-Haddad A, Moreh E, Twito D, Tuchner-Arieli M, Meiner Z. Novel virtual reality system integrating online self-face viewing and mirror visual feedback for stroke rehabilitation: rationale and feasibility. *Top Stroke Rehabil.* 2012 Jul-Aug;19(4):277-86.
15. Yang YR, Chen YH, Chang HC, Chan RC, Wei SH, Wang RY. Effects of interactive visual feedback training on post-stroke pusher syndrome: a pilot randomized controlled study. *Clin Rehabil.* 2015 Oct;29(10):987-93.
16. Michielsen ME, Smits M, Ribbers GM, Stam HJ, van der Geest JN, Bussmann JB, Selles RW. J The neuronal correlates of mirror therapy: an fMRI study on mirror induced visual illusions in patients with stroke. *Neurol Neurosurg Psychiatry.* 2011 Apr;82(4):393-8.
17. Mehnert J, Brunetti M, Steinbrink J, Niedeggen M, Dohle C. Effect of a mirror-like illusion on activation in the precuneus assessed with functional near-infrared spectroscopy. *J Biomed Opt.* 2013 Jun;18(6):066001.
18. Wang J, Fritsch C Bernarding J Krause T Mauritz KH, Brunetti M, Dohle C. Cerebral activation evoked by the mirror illusion of the hand in stroke

patients compared to normal subjects.

NeuroRehabilitation. 2013;33(4):593-603

19. Hamzei F, Lappchen CH, Glauche V, Mader I, Rijntjes M, Weiller C. Functional plasticity induced by mirror training: the mirror as the element connecting both hands to one hemisphere. *Neurorehabil Neural Repair*. 2012 Jun;26(5):484-96.
20. Fritzsche C, Wang J, Dos Santos LF, Mauritz KH, Brunetti M, Dohle C. Different effects of the mirror illusion on motor and somatosensory processing. *Restor Neurol Neurosci*. 2014;32(2):269-80.
21. Lee YY, Hsieh YW, Wu CY, Lin KC, Chen CK. Proximal Fugl-Meyer Assessment Scores Predict Clinically Important Upper Limb Improvement After 3 Stroke Rehabilitative Interventions. *Arch Phys Med Rehabil*. 2015 Dec;96(12):2137-44.
22. Lee YY, Lin KC, Wu CY, Liao CH, Lin JC, Chen CL. Combining Afferent Stimulation and Mirror Therapy for Improving Muscular, Sensorimotor, and Daily Functions After Chronic Stroke: A Randomized, Placebo-Controlled Study. *Am J Phys Med Rehabil*. 2015 Oct;94(10 Suppl 1):859-68.
23. Wu CY, Huang PC, Chen YT, Lin KC, Yang HW. Effects of mirror therapy on motor and sensory recovery in chronic stroke: a randomized controlled trial. *Arch Phys Med Rehabil*. 2013 Jun;94(6):1023-30.
24. Kim K, Lee S, Kim D, Lee K, Kim Y. Effects of mirror therapy combined with motor tasks on upper extremity function and activities daily living of stroke patients. *J Phys Ther Sci*. 2016 Jan;28(2):483-7.
25. In TS, Cha YR, Jung JH, Jung KS. Effects of visual feedback with a mirror on balance ability in patients with stroke. *J Phys Ther Sci*. 2016 Jan; 28(1): 181–185.
26. Kim H, Shim J. Investigation of the effects of mirror therapy on the upper extremity functions of stroke patients using the manual function test. *J Phys Ther Sci*. 2015 Jan;27(1):227-9.
27. Cho HS, Cha HG. Effect of mirror therapy with tDCS on functional recovery of the upper extremity of stroke patients. *J Phys Ther Sci*. 2015 Apr;27(4):1045-7.

28. Choi M, Yoo J, Shin S, Lee W. The effects of stepper exercise with visual feedback on strength, walking, and stair climbing in individuals following stroke. *J Phys Ther Sci.* 2015 Jun;27(6):1861-4.
29. Kim JH, Lee BH. Mirror therapy combined with biofeedback functional electrical stimulation for motor recovery of upper extremities after stroke: a pilot randomized controlled trial. *Occup Ther Int.* 2015 Jun;22(2):51-60.
30. Lee D, Lee M, Lee K, Song C. Asymmetric training using virtual reality reflection equipment and the enhancement of upper limb function in stroke patients: a randomized controlled trial. *J Stroke Cerebrovasc Dis.* 2014 Jul;23(6):1319-26.
31. Selles RW, Michielsen ME, Bussmann JB, Stam HJ, Hurkmans HL, Heijnen I, de Groot D, Ribbers GM. Effects of a mirror-induced visual illusion on a reaching task in stroke patients: implications for mirror therapy training. *Neurorehabil Neural Repair.* 2014 Sep;28(7):652-9.
32. Michielsen ME, Selles RW, van der Geest JN, Eckhardt M, Yavuzer G, Stam HJ, Smits M, Ribbers GM, Bussmann JB. Motor recovery and cortical reorganization after mirror therapy in chronic stroke patients: a phase II randomized controlled trial. *Neurorehabil Neural Repair.* 2011 Mar-Apr;25(3):223-33.
33. Arya KN, Pandian S, Kumar D, Puri V. Task-Based Mirror Therapy Augmenting Motor Recovery in Poststroke Hemiparesis: A Randomized Controlled Trial. *J Stroke Cerebrovasc Dis.* 2015 Aug;24(8):1738-48.
34. Arya KN, Pandian S. Effect of task-based mirror therapy on motor recovery of the upper extremity in chronic stroke patients: a pilot study. *Top Stroke Rehabil.* 2013 May-Jun;20(3):210-7.
35. Rodrigues LC, Farias NC, Gomes RP, Michaelsen SM. Feasibility and effectiveness of adding object-related bilateral symmetrical training to mirror therapy in chronic stroke: A randomized controlled pilot study. *Physiother Theory Pract.* 2016 Feb;32(2):83-91.
36. Colomer C, NOè E, Llorens R. Mirror therapy in chronic stroke survivors with severely impaired upper limb function: a randomized controlled trial. *Eur J Phys Rehabil Med.* 2016 Feb 29.

37. Ji SG, Kim MK. The effects of mirror therapy on the gait of subacute stroke patients: a randomized controlled trial. *Clin Rehabil.* 2015 Apr;29(4):348-54.
38. Kim H, Lee G, Song C. Effect of functional electrical stimulation with mirror therapy on upper extremity motor function in poststroke patients. *J Stroke Cerebrovasc Dis.* 2014 Apr;23(4):655-61.
39. Yoon JA, Koo BI, Shin MJ, Shin YB, Ko HY, Shin YI. Effect of constraint-induced movement therapy and mirror therapy for patients with subacute stroke. *Ann Rehabil Med.* 2014 Aug;38(4):458-66.
40. Lee MM, Cho HY, Song CH. The mirror therapy program enhances upper-limb motor recovery and motor function in acute stroke patients. *Am J Phys Med Rehabil.* 2012 Aug;91(8):689-96, quiz 697-700.
41. Thieme H, Bayn M, Wurg M, Zange C, Pohl M, Behrens J. Mirror therapy for patients with severe arm paresis after stroke--a randomized controlled trial. *Clin Rehabil.* 2013 Apr;27(4):314-24.
42. Invernizzi M, Negrini S, Carda S, Lanzotti L, Cisari C, Baricich A. The value of adding mirror therapy for upper limb motor recovery of subacute stroke patients: a randomized controlled trial. *Eur J Phys Rehabil Med.* 2013 Jun;49(3):311-7.
43. Radajewska A, Opara JA, Kucio C, Błaszczyszyn M, Mehlich K, Szczygiel J. The effects of mirror therapy on arm and hand function in subacute stroke in patients. *Int J Rehabil Res.* 2013 Sep;36(3):268-74.
44. Samuelkamaleshkumar S, Reethajanetsureka S, Pauljebaraj P, Benshamir B, Padankatti SM, David JA. Mirror therapy enhances motor performance in the paretic upper limb after stroke: a pilot randomized controlled trial. *Arch Phys Med Rehabil.* 2014 Nov;95(11):2000-5.
45. Pandian JD, Arora R, Kaur P, Sharma D, Vishwambaran DK, Arima H. Mirror therapy in unilateral neglect after stroke (MUST trial): a randomized controlled trial. *Neurology.* 2014 Sep 9;83(11):1012-7.
46. Mirela Cristina L, Matei D, Ignat B, Popescu CD. Mirror therapy enhances upper extremity motor recovery in stroke patients. *Acta Neurol Belg.* 2015 Dec;115(4):597-603.

47. Brunetti M, Morkisch N, Fritzsich C, Mehnert J, Steinbrink J, Niedeggen M, Dohle C. Potential determinants of efficacy of mirror therapy in stroke patients--A pilot study. *Restor Neurol Neurosci*. 2015;33(4):421-34.
48. Radajewska A, Opara J, Biliński G, Kaczorowska A, Nawrat-Szołtysik A, Kucińska A, Lepsy E. Effectiveness of Mirror Therapy for Subacute Stroke in Relation to Chosen Factors. *Rehabil Nurs*. 2016 Apr 14.
49. Yun GJ, Chun MH, Park JY, Kim BR. The synergic effects of mirror therapy and neuromuscular electrical stimulation for hand function in stroke patients. *Ann Rehabil Med*. 2011 Jun;35(3):316-21.
50. Mohan U, Babu SK, Kumar KV, Suresh BV, Misri ZK, Chakrapani M. Effectiveness of mirror therapy on lower extremity motor recovery, balance and mobility in patients with acute stroke: A randomized sham-controlled pilot trial. *Ann Indian Acad Neurol*. 2013 Oct;16(4):634-9.
51. Yeldan I, Huseyinsinoglu BE, Akinci B, Tarakci E, Baybas S, Ozdincler AR. The effects of very early mirror therapy on functional improvement of the upper extremity in acute stroke patients. *J Phys Ther Sci*. 2015 Nov;27(11):3519-24.
52. Lee HM, Li PC, Fan SC. Delayed mirror visual feedback presented using a novel mirror therapy system enhances cortical activation in healthy adults. *J Neuroeng Rehabil*. 2015 Jul 11;12:56.
53. Zult T, Goodall S, Thomas K, Solnik S, Hortobágyi T, Howatson G. Mirror Training Augments the Cross-education of Strength and Affects Inhibitory Paths. *Med Sci Sports Exerc*. 2016 Jun;48(6):1001-13.
54. Byblow WD, Stinear CM, Smith MC, Bjerre L, Florko BK, McCambridge AB. Mirror symmetric bimanual movement priming can increase corticomotor excitability and enhance motor learning. *PLoS One*. 2012;7(3):e33882.
55. de Almeida Oliveira R, Cintia Dos Santos Vieira P, Rodrigues Martinho Fernandes LF, Patrizzi LJ, Ferreira de Oliveira S, Pascucci Sande de Souza LA. Mental practice and mirror therapy associated with conventional physical therapy training on the hemiparetic upper limb in poststroke rehabilitation: a preliminary study. *Top Stroke Rehabil*. 2014 Nov-Dec;21(6):484-94.

56. Cusmano I, Sterpi I, Mazzone A, Ramat S, Delconte C, Pisano F, Colombo R. Evaluation of upper limb sense of position in healthy individuals and patients after stroke. *J Healthc Eng.* 2014;5(2):145-62.
57. Arya KN, Pandian S, Kumar D, Puri V. Task-based mirror therapy augmenting motor recovery in poststroke hemiparesis: A randomized controlled trial. *J Stroke Cerebrovasc Dis* 2015;24:1738-48.
58. Furukawa K, Suzuki H, Fukuda J. Motion of the drawing hand induces a progressive increase in muscle activity of the non-dominant hand in Ramachandran's mirror-box therapy. *J Rehabil Med* 2012;44:939-43.
59. Michielsen ME, Selles RW, van der Geest JN, Eckhardt M, Yavuzer G, Stam HJ, et al. Motor recovery and cortical reorganization after mirror therapy in chronic stroke patients: A phase II randomized controlled trial. *Neurorehabil Neural Repair* 2011;25:223-33.
60. Rossiter HE, Borrelli MR, Borchert RJ, Bradbury D, Ward NS. Cortical mechanisms of mirror therapy after stroke. *Neurorehabil Neural Repair* 2015;29:444-52.
61. Shumway-Cook A, Woollacott MH. *Motor control: Translating Research into Clinical Practice.* Lippincott Williams and Wilkins; 2007.
62. Stein J, Harvey RL, Macko RF, Winstein CJ, Zorowitz RD. *Stroke recovery and rehabilitation.* New York, NY: Demos Publishing Company; 2009.
63. Harmsen WJ, Bussmann JB, Selles RW, Hurkmans HL, Ribbers GM. A mirror therapy-based action observation protocol to improve motor learning after stroke. *Neurorehabil Neural Repair* 2015;29:509-16.
64. Yavuzer G, Selles R, Sezer N, Sütbeyaz S, Bussmann JB, Köseoğlu F, et al. Mirror therapy improves hand function in subacute stroke: A randomized controlled trial. *Arch Phys Med Rehabil* 2008;89:393-8.
65. Thieme H, Mehrholz J, Pohl M, Behrens J, Dohle C. Mirror therapy for improving motor function after stroke. *Stroke* 2013;44:e1-2.

66. von Rein E, Hoff M, Kaminski E, Sehm B, Steele CJ, Villringer A, et al. Improving motor performance without training: The effect of combining mirror visual feedback with transcranial direct current stimulation. *J Neurophysiol* 2015;113:2383-9.
67. Lee YY, Lin KC, Wu CY, Liao CH, Lin JC, Chen CL. Combining afferent stimulation and mirror therapy for improving muscular, sensorimotor, and daily functions after chronic stroke: A randomized, placebo-controlled study. *Am J Phys Med Rehabil* 2015;94(Suppl 1):859-68.
68. Lin KC, Chen YT, Huang PC, Wu CY, Huang WL, Yang HW, et al. Effect of mirror therapy combined with somatosensory stimulation on motor recovery and daily function in stroke patients: A pilot study. *J Formos Med Assoc* 2014;113:422-8.
69. Sutbeyaz S, Yavuzer G, Sezer N, Koseoglu BF. Mirror therapy enhances lower-extremity motor recovery and motor functioning after stroke: A randomized controlled trial. *Arch Phys Med Rehabil* 2007;88:555-9.
70. Milde C, Rance M, Kirsch P, Trojan J, Fuchs X, Foell J, et al. Do mirror glasses have the same effect on brain activity as a mirror box? Evidence from a functional magnetic resonance imaging study with healthy subjects. *PLoS One* 2015;10:e0127694.
71. Wang J, Fritsch C, Bernarding J, Holtze S, Mauritz KH, Brunetti M, et al. A comparison of neural mechanisms in mirror therapy and movement observation therapy. *J Rehabil Med* 2013;45:410-3.
72. Garry MI, Loftus A, Summers JJ. Mirror, mirror on the wall: Viewing a mirror reflection of unilateral hand movements facilitates ipsilateral M1 excitability. *Exp Brain Res* 2005;163:118-22.
73. Hadoush H, Mano H, Sunagawa T, Nakanishi K, Ochi M. Optimization of mirror therapy to excite ipsilateral primary motor cortex. *NeuroRehabilitation* 2013;32:617-24.

74. Nojima I, Mima T, Koganemaru S, Thabit MN, Fukuyama H, Kawamata T. Human motor plasticity induced by mirror visual feedback. *J Neurosci* 2012;32:1293-300.
75. Fukumura K, Sugawara K, Tanabe S, Ushiba J, Tomita Y. Influence of mirror therapy on human motor cortex. *Int J Neurosci* 2007;117:1039-48.
76. Lappchen CH, Ringer T, Blessin J, Seidel G, Grieshammer S, Lange R, et al. Optical illusion alters M1 excitability after mirror therapy: A TMS study. *J Neurophysiol* 2012;108:2857-61.
77. Avanzino L, Raffo A, Pelosin E, Ogliastro C, Marchese R, Ruggeri P, et al. Training based on mirror visual feedback influences transcallosal communication. *Eur J Neurosci* 2014;40:2581-8.
78. Kang YJ, Ku J, Kim HJ, Park HK. Facilitation of corticospinal excitability according to motor imagery and mirror therapy in healthy subjects and stroke patients. *Ann Rehabil Med* 2011;35:747-58.
79. Deconinck FJ, Smorenburg AR, Benham A, Ledebt A, Feltham MG, Savelsbergh GJ. Reflections on mirror therapy: A systematic review of the effect of mirror visual feedback on the brain. *Neurorehabil Neural Repair* 2015;29:349-61.
80. Bhasin A, Padma Srivastava MV, Kumaran SS, Bhatia R, Mohanty S. Neural interface of mirror therapy in chronic stroke patients: A functional magnetic resonance imaging study. *Neurol India* 2012;60:570-6.
81. Saleh S, Adamovich SV, Tunik E. Mirrored feedback in chronic stroke: Recruitment and effective connectivity of ipsilesional sensorimotor networks. *Neurorehabil Neural Repair* 2014;28:344-54.
82. Carvalho D, Teixeira S, Lucas M, Yuan TF, Chaves F, Peressutti C, et al. The mirror neuron system in post-stroke rehabilitation. *Int Arch Med* 2013;6:41.
83. Mehta UM, Waghmare AV, Thirthalli J, Venkatasubramanian G, Gangadhar BN. Is the human mirror neuron system plastic? Evidence from a transcranial magnetic stimulation study. *Asian J Psychiatr* 2015;17:71-7.

84. Arya KN, Pandian S, Kumar D. Task-based mirror therapy enhances ipsilesional motor functions in stroke: A pilot study. *J Bodyw Mov Ther.* 2017 Apr;21(2):334-341. doi: 10.1016/j.jbmt.2016.08.001. Epub 2016 Aug 11. PubMed PMID: 28532877.
85. Pérez-Cruzado, D., Merchán-Baeza, J. A., González-Sánchez, M. and Cuesta-Vargas, A. I. (2017), Systematic review of mirror therapy compared with conventional rehabilitation in upper extremity function in stroke survivors. *Aust Occup Ther J*, 64: 91–112. doi:10.1111/1440-1630.12342
86. Schick T, Schlake HP, Kallusky J, Hohlfeld G, Steinmetz M, Tripp F, Krakow K, Pinter M, Dohle C. (2017) Synergy effects of combined multichannel EMG-triggered electrical stimulation and mirror therapy in subacute stroke patients with severe or very severe arm/hand paresis. *Restor Neurol Neurosci.* 2017 May 9. doi:10.3233/RNN-160710. [Epub ahead of print] PubMed PMID: 28506003.

## **Mirror Visual Feedback for Conditions other than Stroke:**

### **Phantom Limbs**

The use of visual feedback to treat neurological dysfunction and pain was first introduced by Dr. Ramachandran UCSD.

1. Ramachandran VS, Hirstein W. The perception of phantom limbs. The D. O. Hebb lecture. *Brain* 1998;121:1603-30.  
<https://www.ncbi.nlm.nih.gov/pubmed/9762952>
2. Ramachandran VS, Rogers-Ramachandran D, Cobb S. Touching the phantom limb. *Nature* 1995;377:489-90.
3. Kim SY, Kim YY. Mirror therapy for phantom limb pain. *Korean J Pain.* 2012 Oct;25(4):272-4.
4. Weeks SR, Tsao JW. Incorporation of another person's limb into body image relieves phantom limb pain: a case study. *Neurocase.* 2010 Dec;16(6):461-5.

5. Chan BL, Witt R, Charrow AP, Magee A, Howard R, Pasquina PF, et al . Mirror therapy for phantom limb pain. N Engl J Med 2007;357:2206-7.

### **RSD (CRPS)**

1. McCabe CS, Haigh RC, Ring EF, Halligan PW, Wall PD, Blake DR. A controlled pilot study of the utility of mirror visual feedback in the treatment of complex regional pain syndrome (type 1). Rheumatology (Oxford) 2003b;42:97-101.
2. Cacchio A, De Blasis E, Necozone S, di Orio F, Santilli V. Mirror therapy for chronic complex regional pain syndrome type 1 and stroke. N Engl J Med. 2009 Aug 6;361(6):634-6.
3. Pervane Vural S, Nakipoglu Yuzer GF, Sezgin Ozcan D, Demir Ozbudak S, Ozgirgin N. Effects of Mirror Therapy in Stroke Patients With Complex Regional Pain Syndrome Type 1: A Randomized Controlled Study. Arch Phys Med Rehabil. 2016 Apr;97(4):575-81.
4. Dominique Mouraux , Eric Brassinne , Stéphane Sobczak , Antoine Nonclercq, Warzée , Phillip S. Sizer , Turgay Tuna & Benoît Penelle (2017) 3D augmented reality mirror visual feedback therapy applied to the treatment of persistent, unilateral upper extremity neuropathic pain: a preliminary study, Journal of Manual & Manipulative Therapy, 25:3, 137-143, DOI: 10.1080/10669817.2016.1176726  
<http://dx.doi.org/10.1080/10669817.2016.1176726>

### **Hand Therapy**

1. Yavuzer G, Selles R, Sezer N, Sütbeyaz S, Bussmann JB, Köseoğlu F, et al . Mirror therapy improves hand function in subacute stroke: a randomized controlled trial. Arch Phys Med Rehabil 2008;89:393-8.
2. Altschuler EL, Hu J. Mirror therapy in a patient with a fractured wrist and no active wrist extension. Scand J Plast Reconstr Surg Hand Surg 2008;42:110-1.

3. Rosén B, Lundborg G. Training with a mirror in rehabilitation of the hand. *Scand J Plast Reconstr Surg Hand Surg* 2005;39:104-8.
4. Rostami HR, Arefi A, Tabatabaei S. Effect of mirror therapy on hand function in patients with hand orthopaedic injuries: a randomized controlled trial. *Disabil Rehabil.* 2013 Sep;35(19):1647-51.

### **Neglect syndrome**

1. Ng MJ, Singh P, Pandian JD, Arora R, Kaur P. Mirror therapy in unilateral neglect after stroke (MUST trial): a randomized controlled trial. *Neurology.* 2015 Mar 24;84(12):1286.

### **Alien Hand Syndrome**

1. Romano D, Sedda A, Dell'aquila R, Dalla Costa D, Beretta G, Maravita A, Bottini G. Controlling the alien hand through the mirror box. A single case study of alien hand syndrome. *Neurocase.* 2014 Jun;20(3):307-16.

### **Corpus Callosum Disconnection**

1. Nojima I, Oga T, Fukuyama H, Kawamata T, Mima T. Mirror visual feedback can induce motor learning in patients with callosal disconnection. *Exp Brain Res.* 2013 May;227(1):79-83.

### **Mirror Visual Feedback for Neuro-typicals**

1. Holmes NP, Crozier G, Spence C. When mirrors lie: 'visual capture' of arm position impairs reaching performance. *Cogn Aff Behav Neurosci* 2004;4:193-200.
2. Holmes NP, Snijders HJ, Spence C. Reaching with alien limbs: Visual exposure to prosthetic hands in a mirror biases proprioception without accompanying illusions of ownership. *Percept Psychophys* 2006;68:685-701.
3. Holmes NP, Spence C. Visual bias of unseen hand position with a mirror: spatial and temporal factors. *Exp Brain Res* 2005;166:489-97.

4. Diers M, Kamping S, Kirsch P, Rance M, Bekrater-Bodmann R, Foell J, Trojan J, Fuchs X, Bach F, Maaß H, Cakmak H, Flor H. Illusion-related brain activations: a new virtual reality mirror box system for use during functional magnetic resonance imaging. *Brain Res.* 2015 Jan 12;1594:173-82.
5. Guo F, Xu Q, Abo Salem HM, Yao Y, Lou J, Huang X. The neuronal correlates of mirror therapy: A functional magnetic resonance imaging study on mirror-induced visual illusions of ankle movements. *Brain Res.* 2016 May 15;1639:186-93.
6. Numata K, Murayama T, Takasugi J, Monma M, Oga M. Mirror observation of finger action enhances activity in anterior intraparietal sulcus: a functional magnetic resonance imaging study. *J Jpn Phys Ther Assoc.* 2013;16(1):1-6.
7. Sehm B, Perez MA, Xu B, Hidler J, Cohen LG. Functional neuroanatomy of mirroring during a unimanual force generation task. *Cereb Cortex.* 2010 Jan;20(1):34-45.
8. Bartur G, Pratt H, Dickstein R, Frenkel-Toledo S, Geva A, Soroker N. Electrophysiological manifestations of mirror visual feedback during manual movement. *Brain Res* 2015;1606:113-24.
9. Tominaga W, Matsubayashi J, Deguchi Y, Minami C, Kinai T, Nakamura M, et al. A mirror reflection of a hand modulates stimulus-induced 20-Hz activity. *Neuroimage* 2009;46:500-4.
10. Matthys K, Smits M, Van der Geest JN, Van der Lugt A, Seurinck R, Stam HJ, et al. Mirror-induced visual illusion of hand movements: A functional magnetic resonance imaging study. *Arch Phys Med Rehabil* 2009;90:675-81.
11. Butorina A, Prokofyev A, Nazarova M, Litvak V, Stroganova T. The mirror illusion induces high gamma oscillations in the absence of movement. *Neuroimage* 2014;103:181-91.
12. Fritsch C, Wang J, Dos Santos LF, Mauritz KH, Brunetti M, Dohle C. Different effects of the mirror illusion on motor and somatosensory processing. *Restor Neurol Neurosci* 2014;32:269-80.

### **Back Pain**

1. Wand BM, Tulloch VM, George PJ, Smith AJ, Goucke R, O'Connell NE, Moseley GL. Seeing it helps: movement-related back pain is reduced by

visualization of the back during movement. Clin J Pain. 2012;28(7):602–608. doi: 10.1097/AJP.0b013e31823d480c.

### **Trigeminal Neuralgia**

1. Annegret Hagenberg Clear Effect of Mirror Therapy on Trigeminal Neuralgia Pain(2014). Journal of Observational Pain Medicine – Volume 1, Number 3. ISSN 2047-0800
2. Dabgotra M, Das G, Malik S, Bhattarai R. Mirror Visual Feedback Treatment for Meralgia Paresthetica. J Recent Adv Pain 2016;2(2):59-61.

### **Parkinson's**

1. Gaia Bonassi, Elisa Pelosin, Carla Ogliaastro, Cecilia Cerulli, Giovanni Abbruzzese, Laura Avanzino(2016). Mirror Visual Feedback to Improve Bradykinesia in Parkinson's Disease. Neural Plast 2016 1;2016:8764238. Epub 2016 Aug 1.

### **Other Related Articles**

1. Pandian S, Arya KN. Motor impairment of the ipsilesional body side in poststroke subjects. J Bodyw Mov Ther 2013;17:495-503.
2. Pandian S, Arya KN, Kumar D. Effect of motor training involving the less-affected side (MTLA) in post-stroke subjects: A pilot randomized controlled trial. Top Stroke Rehabil 2015;22:357-67.
3. Louw A, Puentedura EJ, Reese D, Parker P, Miller T, Mintken P. Immediate effects of mirror therapy in patients with shoulder pain and decreased range of motion. Arch Phys Med Rehabil. 2017 May 5. pii: S0003-9993(17)30301-5. doi: 10.1016/j.apmr.2017.03.031. [Epub ahead of print] PubMed PMID: 28483657.

## **Abstracts of Recent Review Articles on Mirror Visual Feedback for Stroke**

### **1. UNDERLYING NEURAL MECHANISMS OF MIRROR THERAPY: IMPLICATIONS FOR MOTOR REHABILITATION IN STROKE**

Kamal Narayan Arya

Department of Occupational Therapy, Pt. Deendayal Upadhyaya Institute for the Physically Handicapped, New Delhi, India

### **Abstract**

Mirror therapy (MT) is a valuable method for enhancing motor recovery in poststroke hemiparesis. The technique utilizes the mirror-illusion created by the movement of sound limb that is perceived as the paretic limb. MT is a simple and economical technique than can stimulate the brain noninvasively. The intervention unquestionably has neural foundation. But the underlying neural mechanisms inducing motor recovery are still unclear. In this review, the neural-modulation due to MT has been explored. Multiple areas of the brain such as the occipital lobe, dorsal frontal area and corpus callosum are involved during the simple MT regime. Bilateral premotor cortex, primary motor cortex, primary somatosensory cortex, and cerebellum also get reorganized to enhance the function of the damaged brain. The motor areas of the lesioned hemisphere receive visuo-motor processing information through the parieto-occipital lobe. The damaged motor cortex responds variably to the MT and may augment true motor recovery. Mirror neurons may also play a possible role in the cortico-stimulatory mechanisms occurring due to the MT.

### **Citation**

Arya KN. Underlying neural mechanisms of mirror therapy: Implications for motor rehabilitation in stroke. *Neurol India* 2016;64:38-44

## **2. MIRROR THERAPY COMBINED WITH BIOFEEDBACK FUNCTIONAL ELECTRICAL STIMULATION FOR MOTOR RECOVERY OF UPPER EXTREMITIES AFTER STROKE: A PILOT RANDOMIZED CONTROLLED TRIAL**

Jung Hee Kim Byoung, Hee Lee

### **Abstract**

The objective of this study was to evaluate the effects of mirror therapy in combination with biofeedback functional electrical stimulation (BF-FES) on motor recovery of the upper extremities after stroke. Twenty-nine patients who suffered a stroke > 6 months prior participated in this study and were randomly allocated to three groups. The BF-FES + mirror therapy and FES + mirror therapy groups practiced training for 5 × 30min sessions over a 4-week period. The control group received a conventional physical therapy program. The following clinical tools were used to assess motor recovery of the upper extremities: electrical muscle tester, electrogoniometer, dual-inclinometer, electrodynamometer, the Box and Block Test (BBT) and Jabsen Taylor Hand Function Test (JHFT), the Functional Independence Measure, the Modified Ashworth Scale, and the Stroke Specific Quality of Life

(SSQOL) assessment. The BF-FES + mirror therapy group showed significant improvement in wrist extension as revealed by the Manual Muscle Test and Range of Motion ( $p < 0.05$ ). The BF-FES + mirror therapy group showed significant improvement in the BBT, JTHT, and SSQOL compared with the FES + mirror therapy group and control group ( $p < 0.05$ ). We found that BF-FES + mirror therapy induced motor recovery and improved quality of life. These results suggest that mirror therapy, in combination with BF-FES, is feasible and effective for motor recovery of the upper extremities after stroke. Copyright © 2014 John Wiley & Sons, Ltd.

### **Citation**

Kim, J. H., & Lee, B. (2014). Mirror Therapy Combined With Biofeedback Functional Electrical Stimulation for Motor Recovery of Upper Extremities After Stroke: A Pilot Randomized Controlled Trial. *Occupational Therapy International*, 22(2), 51-60. doi:10.1002/oti.1384

## **MVF for Other Disorders**

### **1. EFFECT OF MIRROR VISUAL FEEDBACK ON HAND FUNCTIONS IN CHILDREN WITH HEMIPARESIS**

Mohammed Ismael Elsepae, Eman Ibrahim Elhadidy, Hatem Abd Al-Mohsen Emar, Elham Abd Elghaffar Nawar

#### **Background**

Hemiparetic and hemiplegic cerebral palsy (CP) constitute at least a third of all people with CP. Children with hemiparesis are suffering from weak hand muscles and retarded hand use. Mirror therapy is a relatively new approach in rehabilitation used in different neurological disorders. In mirror therapy a mirror is positioned orthogonally in front of the center of the patient's body. The less-affected (healthy) extremity is moved and observed in the mirror. The purpose of this study was to determine the effect of mirror visual feedback on improving hand functions in children with hemiparesis.

Methods: Forty children with hemiparesis of both sexes, ranged in age from five to seven years old, participated in this study. They were divided randomly into two groups of equal number (control and study). The control group received a specially designed physical therapy exercise program for four successive weeks while the study group received mirror exercise program in addition to the same program of the control group. Hand functions assessments was done using grasping and object manipulation subtests of Peabody developmental motor scale (PDMS-2). Evaluation was performed pre and post treatment program.

#### **Results**

There was no significant difference between both groups in the pre-treatment mean values of all measured variables. Also, the results of this study revealed a significant

improvement in the scores of the PDMS-2 and in grasp strength of the 2 groups. Post treatment results revealed more improvement in favor of the study group as compared with the control group.

### **Conclusion**

Using the mirror visual feedback could help in improving hand functions in children with hemiparesis.

### **Citation**

Elsepaee, M. I. (2016). Effect of Mirror Visual Feedback on Hand Functions in Children with Hemiparesis. *International Journal of Physiotherapy*,3(2). doi:10.15621/ijphy/2016/v3i2/94869

## **2. MIRROR THERAPY FOR COMPLEX REGIONAL PAIN SYNDROME (CRPS)--A LITERATURE REVIEW AND AN ILLUSTRATIVE CASE REPORT**

Samaa Al Sayegh, Tove Filén, Mats Johansson, Susanne Sandström, Gisela Stiewe, Stephen Butler

### **Highlights**

- A patient with long-standing CRPS of the lower extremity, improved with mirror therapy.
- Review of CRPS signs and symptoms and the now accepted diagnostic criteria.
- Review of mirror therapy for pain and rehabilitation of CRPS, phantom limb, and stroke.
- Review and evaluation of the published literature for mirror document effects in good quality studies.

### **Abstract**

Background and purpose

This case of a 42 year old woman with lower extremity Complex Regional Pain Syndrome (CRPS) after a twisting injury of the ankle, effectively treated with the addition of mirror therapy to a rehabilitation programme, prompted a literature review of both CRPS and mirror therapy. Mirror therapy is a newer adjunct to other forms of pain control and functional restoration for treatment of CRPS as well as other difficult clinical problems. This was a required group project as part of a university based course in chronic pain for healthcare workers.

### **Materials and methods**

The PubMed database up to September 26, 2012 was reviewed using four search word groups: "CRPS mirror therapy", "mirror CRPS", "reflex sympathetic dystrophy OR Complex Regional Pain Syndrome AND mirror" and "reflex sympathetic dystrophy OR Complex Regional Pain Syndrome AND mirror + RCT". Nine studies from PubMed met the criteria that this working group had chosen for inclusion in the analysis of mirror therapy as treatment. These references were supplemented

by others on CRPS in order to generate an adequate review of both the syndrome CRPS and mirror therapy itself. Some references were specific for mirror therapy in the treatment of CRPS but others described mirror therapy for the treatment of phantom limb pain, brachial plexus avulsion pain, for physical rehabilitation of stroke related paresis and for rehabilitation after hand surgery.

## **Results**

Criteria for the diagnosis of CRPS including the International Association for the Study of Pain criteria and the Budapest criteria are reviewed with an emphasis on the specificity and sensitivity of the various criteria for clinical and research purposes. The signs and symptoms of CRPS are a part of the criteria review.

The main treatment strategy for CRPS is physical rehabilitation for return of function and mirror therapy is one of many possible strategies to aid in this goal.

The patient in this case report had failed many of the adjunctive therapies and rehabilitation had been unsuccessful until the addition of mirror therapy. She then could progress with physical rehabilitation and return to a more normal life. Mirror therapy techniques are briefly described as part of a discussion of its success with relationship to signs and symptoms as well as to the duration of CRPS (and other syndromes). Some discussion of the theories of the central effects of both CRPS and phantom limb pain and how these are affected by mirror therapy is included.

An analysis of the 9 most relevant articles plus a critique of each is present in table form for review.

## **Conclusions**

There appears to be a clear indication for the use of mirror therapy to be included in the multidisciplinary treatment of CRPS types 1 and 2 with a positive effect on both pain and motor function. There is also evidence that mirror therapy can be helpful in other painful conditions such as post stroke pain and phantom limb pain.

## **Implications**

CRPS is often overlooked as an explanation for obscure pain problems. Prompt diagnosis is essential for effective treatment. Mirror therapy is a newer technique, easy to perform and can be a useful adjunct to aid physical rehabilitation and decrease pain in this population. Much further prospective research on mirror therapy in CRPS is ongoing and is needed to systematize the technique, to clarify the effects and to define the place of this therapy in the multidisciplinary management of CRPS.

## **Citation**

Manen, V. S. (2012). Treatment of Complex Regional Pain Syndrome (CPRS) Type I with Mirror Therapy. *Http://isrctn.org/>*. doi:10.1186/isrctn97144266

See following pages for a summary of our clinical work for non-specialists.



*What phantom limbs and mirrors teach us about the brain* by **Stephanie Hegarty BBC World Service: From the section Magazine comments (5 December 2011).**

---

**In a lab in southern California, scientists are curing the previously incurable with little more than a mirror, and changing our understanding of the brain in the process.**

**Ramachandran's 20-year association with the mirror, and phantom limbs, has driven him to the forefront of experimental neuroscience.**

### **How the mirror works**

**The phantom (or arthritic) hand is placed behind the mirror. When the patient looks into the mirror he feels the reflection of the real hand superimposed on the phantom. He then tries to move both hands.**

**Many patients report they feel the phantom mimicking the movement of the real hand.**

**When the real hand opens its fingers, it looks as though the phantom has opened, and pain is relieved. By doing this repeatedly some patients find the phantom disappears. Providing a visual substitute for the phantom limb effectively "amputates" it.**

**The syndrome occurs in at least 90% of amputees - in two-thirds of those it manifests as an insatiable itch in the missing limb, many feel extreme discomfort or even chronic pain.**

**In most cases, pain-killers and surgical treatment have no effect.**

**Ramachandran's first phantom limb patient - who he calls Victor - lost his arm crossing the Mexican border into the US. He had an itch in his missing hand.**

**When Ramachandran prodded him in the left cheek with a cotton bud, Victor claimed he felt it in his missing left thumb - when he touched his upper lip, Victor though he was prodding his index finger.**

**The neurons that detect sensation in the missing hand, at a loss for anything to do, had somehow started detecting sensation in the face.**

**In this case there was a simple and effective treatment for the itch - scratch the face. But to Ramachandran it also had theoretical implications. It appeared to demonstrate the plasticity of brain modules - their ability to adapt to each other and their environment. This was a radical idea as the established notion at the time was that the brain is made up of independent modules, insulated from each other and hardwired to a specific function. The notion of plasticity was something only a small group of scientists were considering.**

### **The “Mirror-Neuron”**

**In 1994, Ramachandran proved the theory by mapping the brain activity of a group of amputees. Using a magnetic scanner he showed that neuron activity was indeed migrating from the hand area to the face. It was a ground-breaking study.**

**But he believed much more could be gleaned from studying phantom limbs.**

**Since 1999 Ramachandran has been exploring the connections between brain modules through a condition called synaesthesia.**

**Synaesthetes experience a blending of their sensations - they taste letters, hear shapes or see numbers as colours. It affects 2-4% of people, is hereditary and eight times more common among artists and creative people.**

**Ramachandran suggests that synaesthetes have a higher number of connections between modules in the brain that process sensations such as colour and sound. He believes the condition holds clues to unravelling the mysteries of human capacity for language, creativity and abstract thought.**

**Do you have synaesthesia? Try and pick out the twos in this sea of fives. Number-colour synaesthetes will see the twos pop out while normal people have to look at each figure individually to tell them apart.**

**In the mid-1990s he followed the work of Italian scientist Giacomo Rizzolatti, who discovered an entirely new type of neuron that he called the mirror neuron.**

**Rizzolatti observed that certain neurons in the brain of a macaque monkey fired when the monkey reached out and when it watched another monkey reach out. Mirror neurons were later discovered in humans too.**

**Ramachandran began to apply this finding to his work with phantom limbs. If mirror neurons fired when an individual watched someone moving a limb, he conjectured, then visual perception might play an important role in creating the sensation of movement.**

**His next subject, Jimmy, felt that his phantom hand was always agonisingly clenched, with his phantom fingernails digging into his missing hand.**

**Ramachandran put a mirror between Jimmy's arms and asked him to move both his phantom and healthy limb simultaneously, while looking at the reflection of the healthy limb - effectively fooling Jimmy's brain into thinking his phantom was moving in a normal way.**

**Jimmy felt his clenched fist release almost immediately.**

**"This is because you are creating intense sensory conflict - the vision is telling you the limb is moving," Ramachandran explains.**

**"One way the brain deals with conflict is to say, 'To hell with it! There is no arm,' and the arm disappears.**

**"I tell my medical colleagues that it is the first example in the history of medicine of successful amputation of a phantom limb."**

**He called the treatment Mirror Visual Feedback therapy or MVF. But it wasn't until much later that MVF was properly acknowledged by clinicians.**

## **Empathy**

**In 2007, an army medic in the US Dr Jack Tsao, performed a controlled test on 22 amputees with remarkable results. All those using the mirror reported a reduction in pain over four weeks, those using a control reported no result or increased pain.**

**At the UK army's rehabilitation centre, Hedley Court, mirror therapy has also been used for the past four years to help amputee soldiers to manage phantom pain.**

## **Mirror Neurons and Landmines**

**Thiet Nguyen Xuan from Vietnam was hunting for scrap metal on a beach in September when a landmine exploded, blowing off both of his legs and two fingers from his left hand.**

**Since the accident, he has felt a burning sensation in his missing fingers. The End the Pain project has provided him with a mirror and training on how to use it. During the training he reported instant relief in his phantom fingers. Over the next two months the project will monitor his progress.**

**End the Pain hopes to extend mirror therapy to the estimated 300,000 amputees in Vietnam.**

## **End the Pain Project**

**"Prosthesis-wearing is key," says army physiotherapist Major Pete LeFeuvre. Those who wait longer for a fake limb seem to suffer more from phantom pain. This suggests it is the visual feedback of seeing an arm rather than the feedback from nerves within it that stops the brain getting confused.**

**In Vietnam, a project called End the Pain has been running for the past three years to spread the therapy among victims of landmines and leprosy sufferers. It has reached over 100 medical practitioners so far and has also extended the project to amputees in Cambodia and Rwanda.**

**The simple therapy has proven useful with other syndromes that have perplexed doctors such as Complex Regional Pain Syndrome, a term for unexplained pain. And at the Royal National Hospital for Rheumatic Diseases in the UK, an associate of Ramachandran, Professor Candy McCabe is testing the use of mirrors with acute stroke victims.**

**Though it is in its very early stages, the experiment into arthritis at UCSD could provide the broadest use of mirror therapy yet.**

**Much of Ramachandran's work since developing mirror therapy has focused on mirror neurons. He believes these neurons help us understand not only what is happening to our own body, but also to others. They are the basis of empathy, he suggests, our ability to feel what others feel.**

**In 2009 he used the phantom limb again to provide evidence for this theory, showing that sufferers could experience relief from phantom pain merely by watching someone else massaging or flexing their own hand.**

**While others spend million on machines with complicated acronyms, the beauty of Ramachandran's work is that he uses ordinary items such as mirrors, pens and paper.**